HEMORRHOIDS

What are Hemorrhoids?

- Hemorrhoids are swollen veins around the anus.
- Each of us has veins around the anus that tend to stretch under pressure. When these veins swell and bulge, they are called hemorrhoids (or piles).
 - One set of veins is inside the rectum (internal), and another is under the skin around the anus (external).



Hemorrhoids are swollen veins around the anus. They can be internal or external.

- Hemorrhoidal tissue is thought to be helpful in holding back stool when we are at rest and in cushioning muscles when we move our bowels.
- Swelling can be caused by straining to move your bowels, sitting too long on the toilet or other causes, such as pregnancy, obesity or liver disease.



The information provided by the AGA Institute is not medical advice and should not be considered a replacement for seeing a medical professional.

How Common Are Hemorrhoids?

- Hemorrhoids are very common in men and women.
- About half of all people have noticeable hemorrhoids by the age of 50.
- Many people have occasional bleeding from hemorrhoids.
- Women's problems may start during pregnancy.
 - The pressure of the fetus in the abdomen, as well as hormonal changes, cause hemorrhoidal veins to swell.
 - These veins also are placed under severe pressure during the birth of the baby. For most women, such hemorrhoids don't last long.





The information provided by the AGA Institute is not medical advice and should not be considered a replacement for seeing a medical professional.

July 2017 © AGA 2017

Page **2** of **2**

HEMORRHOIDS

Symptoms

Internal Hemorrhoids

- Often cause painless rectal bleeding.
- Cause bright red blood in stool, on toilet paper or in the toilet bowl.
- Bleeding happens when the swollen veins are scratched or broken by straining or rubbing.
- Medicines that thin the blood can make bleeding worse.
- An internal hemorrhoid may come through the anus and become swollen and painful.

External Hemorrhoids

- Cause pain and discomfort around the anus.
- Cause painful swelling or a hard lump around the anus.
 - This happens when a blood clot forms.



Did You Know

- When many people think of hemorrhoids, they often speak of external hemorrhoids, but internal hemorrhoids, anal fissures (painful bowel movements due to a small cut in the anus) and other less common anal conditions can all cause bleeding, pain and itching around the anus.
- Talk to your doctor if you don't feel better after trying over-the-counter products.

In General

- Swelling around the anus can cause bleeding and/or itching, which may start a vicious cycle of symptoms.
- Draining mucus or stool residue may cause itching.
 - This can be caused by either too much rubbing or cleaning around the anus or poor cleaning habits with residual stool around the anus.



The information provided by the AGA Institute is not medical advice and should not be considered a replacement for seeing a medical professional.

HEMORRHOIDS

Treatment and Prevention

While hemorrhoids are uncomfortable, they are not often a sign of something serious or dangerous. There are ways to treat hemorrhoids on your own with over-the-counter products.

If your hemorrhoids do not go away or if you see blood in your stool, tell your doctor right away. A doctor may:

- Look at your anus and feel the inside of the rectum with a gloved finger.
- Place a small tool into the rectum to see hemorrhoids. This is called an anoscopy.
- Look at the anal canal and colon with a flexible scope (colonoscopy or sigmoidoscopy).
 - This is needed when internal hemorrhoids cannot be felt with a finger

Treatment

More Fiber

- You may need to increase the fiber in your diet.
- Eat more fresh fruits, leafy veggies, and whole-grain breads and cereals (especially bran).
- Drink six to eight glasses of fluid (not alcohol) each day.





The information provided by the AGA Institute is not medical advice and should not be considered a replacement for seeing a medical professional.

July 2017 © AGA 2017

- Your doctor may suggest that you use a supplement that gives you more fiber and softens the stool.
 - Softer stools make it easier to move your bowels and lessen pressure on the veins.

Good Hygiene

- Cleaning well after you have a bowel movement can help treat and prevent hemorrhoids.
- Bathe the anus gently after each bowel movement using soft, moist toilet paper.
- Avoid a lot of wiping.
- If needed, you can even use a bath or shower instead of wiping.
- After bathing, dry the anus gently with a soft cloth or towel.
- To protect against irritation, cleanse the anus gently and apply a zinc oxide paste (or powder) to the area.
- Medicated suppositories or creams can be found at the drugstore, but are mostly meant for external hemorrhoids.
- You can also try cold packs, a tub bath, warm soaks (a sitz bath) or bed rest to help calm pain or swelling.
- Any of these home treatments may help the symptoms, and no other treatment may be needed. If symptoms do not go away, see your doctor.

Procedural Treatments

In some cases, internal hemorrhoids that have fallen outside of the anus (prolapsed) or that bleed too much must be removed. Your gastroenterologist may be able to treat them during an outpatient visit to the office or to the hospital. A number of methods can be used to remove or reduce the number of hemorrhoids. Talk to your doctor about recovery and which option is best for you.



The information provided by the AGA Institute is not medical advice and should not be considered a replacement for seeing a medical professional.
July 2017



© AGA 2017

- **Endoscopic:** Your doctor will use a long, thin (about the width of your little finger), flexible tube with a tiny camera on the end to look inside and through which to pass tools. Endoscopic methods use freezing, electrical or laser heat, or infrared light to wipe out the hemorrhoidal tissue.
- **Band ligation:** A rubber band is put around the base of the hemorrhoid, which cuts off blood flow, and the hemorrhoid withers away within a few days. This technique is used only for internal hemorrhoids.
- **Sclerotherapy:** With this, a chemical is injected around the vein to shrink the hemorrhoid.
- **Surgical treatment:** Cutting out the hemorrhoids (hemorrhoidectomy) is sometimes suggested, but carries a painful recovery.

Prevention

- Pass your bowel movements as soon as you feel the urge.
- Try not to strain to get the stool out.
- Be active. Move around, walk and exercise to help move the stools through your body.
- Stay regular by eating enough fiber and drinking enough fluid. Adding fiber to your diet helps to make stools that are softer and easier to pass.

Hemorrhoids – What to Know:

- Hemorrhoids are swollen veins found both inside and outside of your anus and rectum.
- Hemorrhoids can be caused by straining when you move your bowels, not cleaning well enough after a bowel movement or sitting on the toilet too long, among other things.
- Hemorrhoids can feel itchy, swollen or may bleed.
- Hemorrhoids are very common and can often be treated with over the counter options.
- Adding more fiber to your diet and making sure you clean well after a bowel movement can help prevent hemorrhoids.



The information provided by the AGA Institute is not medical advice and should not be considered a replacement for seeing a medical professional.

July 2017 © AGA 2017